


SCARD Pool report for 01-01-2000 to 04-02-2026

Participants	Doctors	1,920	
	Patients	624,615	
Specimens	New lesions	1,722,400	84.29%
	Previously biopsied lesions	320,924	15.71%
	Total lesions	2,043,324	100.00%

Breakdown of new versus previously biopsied specimens


Percentage of new lesions tested which were malignant	62.01%
Percentage of Definitively excised lesions that were malignant	79.31%
Lesions tested to find one melanoma (NNT)	4.64
Percentage of lesions tested for NMSC which were NMSC	76.56%
Ratio of New BCCs : New Melanomas	9:1

Accuracy
Diagnostic sensitivity

Melanomas	74.20% of 55,487
All NMSC	95.28% of 978,035
BCCs	90.59% of 526,368
SCCs	81.63% of 446,560

Positive predictive value


Melanomas	42.31% of 97,304
All NMSC	82.92% of 1,123,800
BCCs	42.43% of 1,123,800
SCCs	65.69% of 554,918

Adequacy
Percentage of definitively managed malignant cases where surgical margins were adequate

BCC	87.66% of 360,421
IEC/Bowens disease	83.48% of 109,584
SCC	87.64% of 131,020
Keratoacanthoma	92.45% of 19,154
Melanoma - in situ	77.03% of 36,266
Melanoma - invasive	75.72% of 10,064
Melanoma - invasive > 1mm	44.59% of 1,350
Other malignant	68.71% of 1,582

Lesion Breakdown
Histological Diagnosis

BCC (unspecified type)	46,294	2.68%
BCC - Superficial	152,438	8.84%
BCC - Nodular/Solid	258,441	14.99%
BCC - Aggressive	69,195	4.01%
IEC/Bowens disease	238,504	13.83%
SCC	181,307	10.51%
Keratoacanthoma	26,749	1.55%
Pinkus Fibroepithelioma	205	0.01%
Merkel cell tumour	194	0.01%
Other malignant	2,381	0.14%
NMSC Metastasis	65	0.00%
Melanoma - in situ	39,920	2.31%
Melanoma - invasive	12,143	0.70%
Melanoma - invasive > 1mm	2,960	0.17%
Melanoma - metastasis	464	0.03%
MELTUMP	2,262	0.13%
Naevus - other	93,716	5.43%
Naevus - dysplastic/Clark	59,423	3.45%
Naevus - blue	6,072	0.35%
Naevus - Spitz/Reed	2,160	0.13%
Naevus - Compound	7,324	0.42%
Solar keratosis	156,387	9.07%
Solar lentigo	23,134	1.34%
Seborrhoeic keratosis	91,048	5.28%
Lentigo Simplex	484	0.03%
Lichenoid keratosis (LPLK)	30,905	1.79%
Dermatofibroma	12,593	0.73%
Sebaceous gland hyperplasia	3,593	0.21%
Benign cyst	26,932	1.56%
Other benign	115,376	6.69%
Histology Pending	2,237	0.13%



BCC - 31.65%
IEC - 14.34%
SCC - 10.90%
Keratoacanthoma - 1.61%
Other malignant - 0.18%
Melanoma - 3.47%
Naevus - 10.14%
Solar keratosis - 9.40%
Solar lentigo - 1.42%
Seborrhoeic keratosis - 5.48%
Lichenoid keratosis - 1.86%
Dermatofibroma - 0.76%
Sebaceous gland - 0.22%
Benign cyst - 1.62%
Other benign - 6.94%

Procedures
Definitive Surgical Management used to exclude melanoma

Ellipse	53,082	85.75%
Flap	1,100	1.78%
Graft - SSG	83	0.13%
Graft - FTSG	199	0.32%
No Closure	134	0.22%
Shave/Saucerisation	2,719	4.39%
Curettage & Cautery	797	1.29%
Liquid N2 freeze/thaw	42	0.07%
PDT	4	0.01%
Imiquimod	7	0.01%
5 FU cream	15	0.02%
GP referral	2,327	3.76%
Specialist referral	870	1.41%
Other	446	0.72%

Biopsy used to exclude melanoma

Punch - sample	15,951	7.61%
Shave - sample	26,338	12.57%
Incisional	3,066	1.46%
Punch - removal	37,915	18.09%
Shave - removal	49,535	23.64%
Excisional	75,589	36.07%
Curettage	801	0.38%
Other	288	0.14%

Breakdown of definitive management procedures for malignant conditions

Ellipse	579,694	64.60%
Flap	76,388	8.51%
Graft - SSG	5,110	0.57%
Graft - FTSG	19,151	2.13%
No Closure	3,301	0.37%
Shave/Saucerisation	13,768	1.53%
Curettage & Cautery	115,344	12.85%
Liquid N2 freeze/thaw	11,804	1.32%
PDT	1,713	0.19%
Imiquimod	8,916	0.99%
5 FU cream	15,624	1.74%
GP referral	12,592	1.40%
Specialist referral	23,224	2.59%
Other	3,824	0.43%

Breakdown of definitive management procedures for benign conditions

Ellipse	87,627	73.72%
Flap	1,540	1.30%
Graft - SSG	67	0.06%
Graft - FTSG	178	0.15%
No Closure	359	0.30%
Shave/Saucerisation	9,740	8.19%
Liquid N2 freeze/thaw	3,785	3.18%
5 FU cream	1,546	1.30%
GP referral	1,209	1.02%
Specialist referral	709	0.60%
Other	2,270	1.91%

Procedures (continued)

Percentage of procedures/closures that were complex

Complex Closures	1.14% of 156,434
Complex Closures (inc Curettage & Cautery)	1.08% of 165,710

Locations

Breakdown of melanomas by location and percentage melanomas of total lesions at that location

Nose	582	0.57% of 102,546
Lip	69	0.25% of 27,488
Ear	1238	1.71% of 72,500
Eyelid	111	0.66% of 16,767
Other face	5441	1.89% of 288,076
Scalp	1386	2.17% of 63,801
Neck	2373	2.68% of 88,538
Shoulder	4033	4.98% of 80,969
Chest	2850	2.79% of 102,242
Abdomen	1279	5.67% of 22,557
Genitalia	21	0.94% of 2,234
Back	17334	6.97% of 248,718
Buttock	161	3.78% of 4,260
Arm	5876	5.76% of 102,091
Forearm	3399	2.60% of 130,801
Hand Dorsal	124	0.19% of 66,381
Hand Palmar	9	1.64% of 550
Finger Dorsal	30	0.23% of 12,829
Finger Nail	20	9.39% of 213
Finger Palmar	3	1.13% of 265
Thigh	2746	5.20% of 52,844
Leg	5751	2.75% of 208,931
Foot Dorsal	371	2.41% of 15,372
Foot Plantar	89	5.48% of 1,624
Toe Dorsal	80	3.32% of 2,410
Toe Nail	7	3.35% of 209
Toe Plantar	13	4.10% of 317
Palm Or Sole	0	0% of 22

