


SCARD Pool report for 01-01-2004 to 31-12-2004

Participants	Doctors	9	
	Patients	181	
Specimens	New lesions	280	85.37%
	Previously biopsied lesions	48	14.63%
	Total lesions	328	100.00%

Breakdown of new versus previously biopsied specimens



Percentage of new lesions tested which were malignant	56.04%
Percentage of Definitively excised lesions that were malignant	62.90%
Lesions tested to find one melanoma (NNT)	4.17
Percentage of lesions tested for NMSC which were NMSC	66.82%
Ratio of New BCCs : New Melanomas	7:1

Accuracy

Diagnostic sensitivity

Melanomas	83.33% of 12
All NMSC	97.87% of 141
BCCs	97.73% of 88
SCCs	79.25% of 53
Malagana	00.010/ -5.11

Positive predictive value

Melanomas	90.91% of 11
All NMSC	92.00% of 150
BCCs	57.33% of 150
SCCs	91.30% of 46

Adequacy


Percentage of definitively managed malignant cases where surgical margins were adequate

BCC	85.94% of 64
IEC/Bowens disease	62.50% of 8
SCC	76.92% of 26
Keratoacanthoma	100.00% of 2
Melanoma - in situ	80.00% of 10
Melanoma - invasive	100.00% of 2
Melanoma - invasive > 1mm	0% of 0
Other malignant	0% of 0



Lesion Breakdown Histological Diagnosis

PCC (unenceified type)	25	8.87%
BCC (unspecified type)	26	9.22%
BCC - Superficial		10.64%
BCC - Nodular/Solid	30 7	
BCC - Aggressive	-	2.48%
IEC/Bowens disease	20	7.09%
SCC	30	10.64%
Keratoacanthoma	3	1.06%
Pinkus Fibroepithelioma	0	0%
Merkel cell tumour	0	0%
Other malignant	0	0%
NMSC Metastasis	0	0%
Melanoma - in situ	10	3.55%
Melanoma - invasive	2	0.71%
Melanoma - invasive > 1mm	0	0%
Melanoma - metastasis	0	0%
MELTUMP	0	0%
Naevus - other	10	3.55%
Naevus - dysplastic/Clark	17	6.03%
Naevus - blue	2	0.71%
Naevus - Spitz/Reed	0	0%
Naevus - Compound	0	0%
Solar keratosis	15	5.32%
Solar lentigo	2	0.71%
Seborrhoeic keratosis	39	13.83%
Lentigo Simplex	0	0%
Lichenoid keratosis (LPLK)	0	0%
Dermatofibroma	0	0%
Sebaceous gland hyperplasia	1	0.35%
Benign cyst	13	4.61%
Other benign	21	7.45%
Histology Pending	2	0.71%
5, 5		





Procedures			
Definitive Surgical Managemen	t used to exclude melanoma		
	Ellipse	38	100.00%
	Flap	0	0%
	Graft - SSG	0	0%
	Graft - FTSG	0	0%
	No Closure	0	0%
	Shave/Saucerisation	0	0%
	Curettage & Cautery	0	0%
	Liquid N2 freeze/thaw	0	0%
	PDT	0	0%
	Imiquimod	0	0%
	5 FU cream	0	0%
	GP referral	0	0%
	Specialist referral	0	0%
	Other	0	0%
Biopsy used to exclude melano	ma		
	Punch - sample	3	25.00%
	Shave - sample	0	0%
	Incisional	0	0%
	Punch - removal	1	8.33%
	Shave - removal	2	16.67%
	Excisional	6	50.00%
	Curettage	0	0%
	Other	0	0%
Breakdown of definitive manag	ement procedures for malignant conditions		
	Ellipse	96	59.63%
	Flap	1	0.62%
	Graft - SSG	0	0%
	Graft - FTSG	0	0%
	No Closure	0	0%
	Shave/Saucerisation	9	5.59%
	Curettage & Cautery	27	16.77%
	Liquid N2 freeze/thaw	1	0.62%
	PDT	0	0%
	Imiquimod	5	3.11%
	5 FU cream	0	0%
	GP referral	0	0%
	Specialist referral	22	13.66%
	Other	0	0%
Breakdown of definitive manag	ement procedures for benign conditions		
	Ellipse	70	88.61%
	Flap	0	0%
	Graft - SSG	0	0%
	Graft - FTSG	0	0%
	No Closure	0	0%
	Shave/Saucerisation	3	3.80%
	Liquid N2 freeze/thaw	0	0%
	5 FU cream	0	0%
	GP referral	0	0%
	Specialist referral	0	0%
	Other	0	0%



Procedures (continued)

Percentage of procedures/closures that were complex

Complex Closures	0.45% of 222
Complex Closures (inc Curettage & Cautery)	0.39% of 255

Locations

Breakdown of melanomas by location and percentage melanomas of total lesions at that location

	incolons at that location	
Nose	0	0% of 10
Lip	0	0% of 3
Ear	0	0% of 7
Eyelid	0	0% of 0
Other face	2	4.65% of 43
Scalp	0	0% of 9
Neck	0	0% of 20
Shoulder	1	7.69% of 13
Chest	1	6.67% of 15
Abdomen	0	0% of 6
Genitalia	0	0% of 0
Back	4	7.84% of 51
Buttock	0	0% of 0
Arm	1	6.25% of 16
Forearm	1	4.55% of 22
Hand Dorsal	0	0% of 10
Hand Palmar	0	0% of 1
Finger Dorsal	0	0% of 6
Finger Nail	0	0% of 0
Finger Palmar	0	0% of 0
Thigh	0	0% of 15
Leg	2	6.90% of 29
Foot Dorsal	0	0% of 3
Foot Plantar	0	0% of 1
Toe Dorsal	0	0% of 0
Toe Nail	0	0% of 0
Toe Plantar	0	0% of 0
Palm Or Sole	0	0% of 0

