


SCARD Pool report for 01-01-2005 to 31-12-2005

Participants	Doctors	11	
	Patients	214	
Specimens	New lesions	353	82.48%
	Previously biopsied lesions	75	17.52%
	Total lesions	428	100.00%

Breakdown of new versus previously biopsied specimens


Percentage of new lesions tested which were malignant	57.51%
Percentage of Definitively excised lesions that were malignant	60.82%
Lesions tested to find one melanoma (NNT)	6.20
Percentage of lesions tested for NMSC which were NMSC	69.49%
Ratio of New BCCs : New Melanomas	12:1

Accuracy
Diagnostic sensitivity

Melanomas	90.00% of 10
All NMSC	98.41% of 189
BCCs	97.50% of 120
SCCs	73.53% of 68

Positive predictive value


Melanomas	75.00% of 12
All NMSC	88.15% of 211
BCCs	55.45% of 211
SCCs	78.13% of 64

Adequacy
Percentage of definitively managed malignant cases where surgical margins were adequate

BCC	94.44% of 54
IEC/Bowens disease	100.00% of 11
SCC	82.76% of 29
Keratoacanthoma	100.00% of 1
Melanoma - in situ	100.00% of 5
Melanoma - invasive	0% of 0
Melanoma - invasive > 1mm	100.00% of 1
Other malignant	0% of 0

Lesion Breakdown
Histological Diagnosis

BCC (unspecified type)	30	8.47%
BCC - Superficial	29	8.19%
BCC - Nodular/Solid	50	14.12%
BCC - Aggressive	11	3.11%
IEC/Bowens disease	25	7.06%
SCC	42	11.86%
Keratoacanthoma	1	0.28%
Pinkus Fibroepithelioma	0	0%
Merkel cell tumour	0	0%
Other malignant	1	0.28%
NMSC Metastasis	0	0%
Melanoma - in situ	6	1.69%
Melanoma - invasive	2	0.56%
Melanoma - invasive > 1mm	2	0.56%
Melanoma - metastasis	0	0%
MELTUMP	0	0%
Naevus - other	13	3.67%
Naevus - dysplastic/Clark	21	5.93%
Naevus - blue	2	0.56%
Naevus - Spitz/Reed	0	0%
Naevus - Compound	0	0%
Solar keratosis	24	6.78%
Solar lentigo	1	0.28%
Seborrhoeic keratosis	42	11.86%
Lentigo Simplex	0	0%
Lichenoid keratosis (LPLK)	0	0%
Dermatofibroma	3	0.85%
Sebaceous gland hyperplasia	1	0.28%
Benign cyst	11	3.11%
Other benign	29	8.19%
Histology Pending	1	0.28%



Procedures
Definitive Surgical Management used to exclude melanoma

Ellipse	37	94.87%
Flap	0	0%
Graft - SSG	0	0%
Graft - FTSG	0	0%
No Closure	0	0%
Shave/Saucerisation	0	0%
Curettage & Cautery	1	2.56%
Liquid N2 freeze/thaw	0	0%
PDT	0	0%
Imiquimod	0	0%
5 FU cream	0	0%
GP referral	0	0%
Specialist referral	1	2.56%
Other	0	0%

Biopsy used to exclude melanoma

Punch - sample	6	25.00%
Shave - sample	5	20.83%
Incisional	0	0%
Punch - removal	1	4.17%
Shave - removal	0	0%
Excisional	12	50.00%
Curettage	0	0%
Other	0	0%

Breakdown of definitive management procedures for malignant conditions

Ellipse	108	56.54%
Flap	3	1.57%
Graft - SSG	0	0%
Graft - FTSG	0	0%
No Closure	0	0%
Shave/Saucerisation	1	0.52%
Curettage & Cautery	19	9.95%
Liquid N2 freeze/thaw	0	0%
PDT	1	0.52%
Imiquimod	17	8.90%
5 FU cream	1	0.52%
GP referral	0	0%
Specialist referral	41	21.47%
Other	0	0%

Breakdown of definitive management procedures for benign conditions

Ellipse	42	67.74%
Flap	0	0%
Graft - SSG	0	0%
Graft - FTSG	0	0%
No Closure	0	0%
Shave/Saucerisation	0	0%
Liquid N2 freeze/thaw	1	1.61%
5 FU cream	2	3.23%
GP referral	0	0%
Specialist referral	1	1.61%
Other	0	0%

Procedures (continued)

Percentage of procedures/closures that were complex

Complex Closures	0%	of 73
Complex Closures (inc Curettage & Cautery)	0%	of 87

Locations

Breakdown of melanomas by location and percentage melanomas of total lesions at that location

Nose	0	0%	of 24
Lip	0	0%	of 1
Ear	0	0%	of 16
Eyelid	0	0%	of 4
Other face	1	1.75%	of 57
Scalp	1	9.09%	of 11
Neck	0	0%	of 15
Shoulder	1	5.88%	of 17
Chest	0	0%	of 22
Abdomen	1	33.33%	of 3
Genitalia	0	0%	of 3
Back	2	3.70%	of 54
Buttock	0	0%	of 0
Arm	2	12.50%	of 16
Forearm	1	4.00%	of 25
Hand Dorsal	0	0%	of 20
Hand Palmar	0	0%	of 0
Finger Dorsal	0	0%	of 3
Finger Nail	0	0%	of 0
Finger Palmar	0	0%	of 0
Thigh	0	0%	of 5
Leg	1	1.96%	of 51
Foot Dorsal	0	0%	of 2
Foot Plantar	0	0%	of 4
Toe Dorsal	0	0%	of 0
Toe Nail	0	0%	of 0
Toe Plantar	0	0%	of 0
Palm Or Sole	0	0%	of 0

