

# SCARD Pool report for 01-01-2005 to 31-12-2005

Participants	Doctors	8	
	Patients	209	
Specimens	New lesions	350	82.74%
-	Previously biopsied lesions	73	17.26%
	Total lesions	423	100.00%

### Breakdown of new versus previously biopsied specimens



Percentage of new lesions tested which were malignant	57.43%
Percentage of Definitively excised lesions that were malignant	60.42%
Lesions tested to find one melanoma (NNT)	6.20
Percentage of lesions tested for NMSC which were NMSC	69.26%
Ratio of New BCCs : New Melanomas	12:1

### **Accuracy**

**Diagnostic sensitivity** 

Melanomas	90.00% of 10
All NMSC	98.40% of 187
BCCs	97.46% of 118
SCCs	73.53% of 68

## Positive predictive value

Melanomas	75.00% of 12
All NMSC	88.04% of 209
BCCs	55.02% of 209
SCCs	78.13% of 64

# **Adequacy**


Percentage of definitively managed malignant cases where surgical margins were adequate

BCC	94.34% of 53
IEC/Bowens disease	100.00% of 11
SCC	82.14% of 28
Keratoacanthoma	100.00% of 1
Melanoma - in situ	100.00% of 5
Melanoma - invasive	0% of 0
Melanoma - invasive > 1mm	100.00% of 1
Other malignant	0% of 0



# Lesion Breakdown Histological Diagnosis

BCC (unspecified type)	30	8.55%
BCC - Superficial	29	8.26%
BCC - Nodular/Solid	49	13.96%
BCC - Aggressive	10	2.85%
IEC/Bowens disease	25	7.12%
SCC	42	11.97%
Keratoacanthoma	1	0.28%
Pinkus Fibroepithelioma	0	0%
Merkel cell tumour	0	0%
Other malignant	1	0.28%
NMSC Metastasis	0	0%
Melanoma - in situ	6	1.71%
Melanoma - invasive	2	0.57%
Melanoma - invasive > 1mm	2	0.57%
Melanoma - metastasis	0	0%
MELTUMP	0	0%
Naevus - other	13	3.70%
Naevus - dysplastic/Clark	21	5.98%
Naevus - blue	2	0.57%
Naevus - Spitz/Reed	0	0%
Naevus - Compound	0	0%
Solar keratosis	24	6.84%
Solar lentigo	1	0.28%
Seborrhoeic keratosis	41	11.68%
Lentigo Simplex	0	0%
Lichenoid keratosis (LPLK)	0	0%
Dermatofibroma	3	0.85%
Sebaceous gland hyperplasia	1	0.28%
Benign cyst	11	3.13%
Other benign	29	8.26%
Histology Pending	1	0.28%





Procedures			
Definitive Surgical Managemen	nt used to exclude melanoma		
3 3	Ellipse	37	94.87%
	Flap	0	0%
	Graft - SSG	0	0%
	Graft - FTSG	0	0%
	No Closure	0	0%
	Shave/Saucerisation	0	0%
	Curettage & Cautery	1	2.56%
	Liquid N2 freeze/thaw	0	0%
	PDT	0	0%
	Imiquimod	0	0%
	5 FU cream	0	0%
	GP referral	0	0%
	Specialist referral	1	2.56%
	Other	0	0%
Biopsy used to exclude meland	oma		
	Punch - sample	6	25.00%
	Shave - sample	5	20.83%
	Incisional	0	0%
	Punch - removal	1	4.17%
	Shave - removal	0	0%
	Excisional	12	50.00%
	Curettage	0	0%
	Other	0	0%
Breakdown of definitive manag	gement procedures for malignant conditions		
	Ellipse	106	56.08%
	Flap	3	1.59%
	Graft - SSG	0	0%
	Graft - FTSG	0	0%
	No Closure	0	0%
	Shave/Saucerisation	1	0.53%
	Curettage & Cautery	19	10.05%
	Liquid N2 freeze/thaw	0	0%
	PDT	1	0.53%
	Imiquimod	17	8.99%
	5 FU cream	1	0.53%
	GP referral	0	0%
	Specialist referral	41	21.69%
	Other	0	0%
Breakdown of definitive manage	gement procedures for benign conditions	42	67.740/
	Ellipse	42	67.74%
	Flap	0	0%
	Graft - SSG	0	0%
	Graft - FTSG	0	0%
	No Closure	0	0%
	Shave/Saucerisation	0	0%
	Liquid N2 freeze/thaw	1	1.61%
	5 FU cream GP referral	2	3.23%
		0	0% 1.61%
	Specialist referral	1	
	Other	0	0%



### **Procedures (continued)**

## Percentage of procedures/closures that were complex

Complex Closures	1.47% of 204
Complex Closures (inc Curettage & Cautery)	1.27% of 237

### Locations

## Breakdown of melanomas by location and percentage melanomas of total lesions at that location

· · · · · · · · · · · · · · · · · · ·		
Nose	0	0% of 24
Lip	0	0% of 1
Ear	0	0% of 16
Eyelid	0	0% of 4
Other face	1	1.79% of 56
Scalp	1	9.09% of 11
Neck	0	0% of 15
Shoulder	1	5.88% of 17
Chest	0	0% of 22
Abdomen	1	33.33% of 3
Genitalia	0	0% of 3
Back	2	3.70% of 54
Buttock	0	0% of 0
Arm	2	12.50% of 16
Forearm	1	4.00% of 25
Hand Dorsal	0	0% of 20
Hand Palmar	0	0% of 0
Finger Dorsal	0	0% of 3
Finger Nail	0	0% of 0
Finger Palmar	0	0% of 0
Thigh	0	0% of 5
Leg	1	2.04% of 49
Foot Dorsal	0	0% of 2
Foot Plantar	0	0% of 4
Toe Dorsal	0	0% of 0
Toe Nail	0	0% of 0
Toe Plantar	0	0% of 0
Palm Or Sole	0	0% of 0

