


## SCARD Pool report for 01-01-2018 to 31-12-2018

| Participants | Doctors<br>Patients         | 426<br>58.247 |         |
|--------------|-----------------------------|---------------|---------|
|              | ratients                    | 30,247        |         |
| Specimens    | New lesions                 | 109,659       | 84.89%  |
|              | Previously biopsied lesions | 19,523        | 15.11%  |
|              | Total lesions               | 129,182       | 100.00% |

#### Breakdown of new versus previously biopsied specimens



| Percentage of new lesions tested which were malignant          | 62.09% |
|--|--------|
| Percentage of Definitively excised lesions that were malignant | 81.39% |
| Lesions tested to find one melanoma (NNT)                      | 4.50   |
| Percentage of lesions tested for NMSC which were NMSC          | 80.60% |
| Ratio of New BCCs : New Melanomas                              | 10:1   |

### **Accuracy**

**Diagnostic sensitivity** 

| Melanomas | 73.70% of 3,422  |
|-----------|------------------|
| All NMSC  | 95.71% of 61,492 |
| BCCs      | 91.91% of 33,176 |
| SCCs      | 82.60% of 27,943 |
|           |                  |

### Positive predictive value

| Melanomas | 43.67% of 5,775  |
|-----------|------------------|
| All NMSC  | 86.61% of 67,954 |
| BCCs      | 44.87% of 67,954 |
| SCCs      | 64.93% of 35,545 |

## **Adequacy**


#### Percentage of definitively managed malignant cases where surgical margins were adequate

| BCC                       | 88.11% of 23,277 |
|---------------------------|------------------|
| IEC/Bowens disease        | 82.42% of 5,101  |
| SCC                       | 87.85% of 8,280  |
| Keratoacanthoma           | 91.53% of 885    |
| Melanoma - in situ        | 77.41% of 2,328  |
| Melanoma - invasive       | 72.77% of 595    |
| Melanoma - invasive > 1mm | 37.35% of 83     |
| Other malignant           | 54.95% of 91     |



# Lesion Breakdown Histological Diagnosis

| DCC (unanacified tune)      | F.C.   | 0.050/ |
|-----------------------------|--------|--------|
| BCC (unspecified type)      | 56     | 0.05%  |
| BCC - Superficial           | 10,162 | 9.26%  |
| BCC - Nodular/Solid         | 17,689 | 16.12% |
| BCC - Aggressive            | 5,269  | 4.80%  |
| IEC/Bowens disease          | 14,397 | 13.12% |
| SCC                         | 12,230 | 11.14% |
| Keratoacanthoma             | 1,316  | 1.20%  |
| Pinkus Fibroepithelioma     | 15     | 0.01%  |
| Merkel cell tumour          | 15     | 0.01%  |
| Other malignant             | 153    | 0.14%  |
| NMSC Metastasis             | 4      | 0.00%  |
| Melanoma - in situ          | 2,503  | 2.28%  |
| Melanoma - invasive         | 688    | 0.63%  |
| Melanoma - invasive > 1mm   | 194    | 0.18%  |
| Melanoma - metastasis       | 37     | 0.03%  |
| MELTUMP                     | 186    | 0.17%  |
| Naevus - other              | 6,117  | 5.57%  |
| Naevus - dysplastic/Clark   | 3,472  | 3.16%  |
| Naevus - blue               | 338    | 0.31%  |
| Naevus - Spitz/Reed         | 125    | 0.11%  |
| Naevus - Compound           | 0      | 0%     |
| Solar keratosis             | 10,352 | 9.43%  |
| Solar lentigo               | 1,313  | 1.20%  |
| Seborrhoeic keratosis       | 5,460  | 4.97%  |
| Lentigo Simplex             | 0      | 0%     |
| Lichenoid keratosis (LPLK)  | 1,984  | 1.81%  |
| Dermatofibroma              | 783    | 0.71%  |
| Sebaceous gland hyperplasia | 233    | 0.21%  |
| Benign cyst                 | 1,809  | 1.65%  |
| Other benign                | 7,328  | 6.68%  |
| Histology Pending           | 107    | 0.10%  |
|                             |        | 0.2070 |





|                       |   | ·       |        |
|-----------------------|---|---------|--------|
| Procedures            |   |         |        |
| Definitive Surgical I | Management used to exclude melanoma           |         |        |
|                       | Ellipse                                       | 2,736   | 83.49% |
|                       | Flap  | 67      | 2.04%  |
|                       | Graft - SSG                                   | 7       | 0.21%  |
|                       | Graft - FTSG                                  | 14      | 0.43%  |
|                       | No Closure                                    | 10      | 0.31%  |
|                       | Shave/Saucerisation                           | 119     | 3.63%  |
|                       | Curettage & Cautery                           | 24      | 0.73%  |
|                       | Liquid N2 freeze/thaw                         | 4       | 0.12%  |
|                       | PDT   | 2       | 0.06%  |
|                       | Imiquimod                                     | 1       | 0.03%  |
|                       | 5 FU cream                                    | 1       | 0.03%  |
|                       | GP referral                                   | 190     | 5.80%  |
|                       | Specialist referral                           | 66      | 2.01%  |
|                       | Other   | 31      | 0.95%  |
| Biopsy used to excl   | ude melanoma                                  |         |        |
|                       | Punch - sample                                | 815     | 6.43%  |
|                       | Shave - sample                                | 1,068   | 8.42%  |
|                       | Incisional                                    | 226     | 1.78%  |
|                       | Punch - removal                               | 2,449   | 19.32% |
|                       | Shave - removal                               | 3,065   | 24.18% |
|                       | Excisional                                    | 5,021   | 39.61% |
|                       | Curettage                                     | 22      | 0.179  |
|                       | Other   | 10      | 0.08%  |
| Breakdown of defin    | itive management procedures for malignant con | ditions |        |
|                       | Ellipse                                       | 34,212  | 61.76% |
|                       | Flap  | 5,261   | 9.50%  |
|                       | Graft - SSG                                   | 360     | 0.65%  |
|                       | Graft - FTSG                                  | 1,335   | 2.41%  |
|                       | No Closure                                    | 306     | 0.55%  |
|                       | Shave/Saucerisation                           | 934     | 1.69%  |
|                       | Curettage & Cautery                           | 6,370   | 11.50% |
|                       | Liquid N2 freeze/thaw                         | 799     | 1.44%  |
|                       | PDT   | 122     | 0.22%  |
|                       | Imiquimod                                     | 571     | 1.03%  |
|                       | 5 FU cream                                    | 1,349   | 2.44%  |
|                       | GP referral                                   | 1,500   | 2.719  |
|                       |   |         |        |

# Breakdown of definitive management procedures for benign conditions

Other

Specialist referral

| nagement procedures for benigh condition | ons   |        |
|--|-------|--------|
| Ellipse                                  | 5,110 | 72.55% |
| Flap                                     | 83    | 1.18%  |
| Graft - SSG                              | 2     | 0.03%  |
| Graft - FTSG                             | 9     | 0.13%  |
| No Closure                               | 36    | 0.51%  |
| Shave/Saucerisation                      | 599   | 8.50%  |
| Liquid N2 freeze/thaw                    | 180   | 2.56%  |
| 5 FU cream                               | 122   | 1.73%  |
| GP referral                              | 165   | 2.34%  |
| Specialist referral                      | 52    | 0.74%  |
| Other                                    | 142   | 2.02%  |

2.56%

0.53%

1,419

293



### **Procedures (continued)**

### Percentage of procedures/closures that were complex

| Complex Closures                           | 10.25% of 43,932 |
|--|------------------|
| Complex Closures (inc Curettage & Cautery) | 9.35% of 48,158  |

### Locations

### Breakdown of melanomas by location and percentage melanomas of total lesions at that location

| ,             |      |                 |
|---------------|------|-----------------|
| Nose          | 35   | 0.54% of 6,521  |
| Lip           | 8    | 0.44% of 1,810  |
| Ear           | 83   | 1.78% of 4,673  |
| Eyelid        | 3    | 0.26% of 1,160  |
| Other face    | 330  | 1.79% of 18,449 |
| Scalp         | 74   | 1.81% of 4,086  |
| Neck          | 161  | 2.98% of 5,406  |
| Shoulder      | 264  | 5.12% of 5,154  |
| Chest         | 174  | 2.70% of 6,443  |
| Abdomen       | 68   | 5.16% of 1,319  |
| Genitalia     | 1    | 0.90% of 111    |
| Back          | 1072 | 6.77% of 15,830 |
| Buttock       | 10   | 3.70% of 270    |
| Arm           | 386  | 5.93% of 6,508  |
| Forearm       | 199  | 2.30% of 8,642  |
| Hand Dorsal   | 7    | 0.15% of 4,704  |
| Hand Palmar   | 0    | 0% of 50        |
| Finger Dorsal | 1    | 0.12% of 867    |
| Finger Nail   | 2    | 12.50% of 16    |
| Finger Palmar | 0    | 0% of 20        |
| Thigh         | 173  | 5.39% of 3,211  |
| Leg           | 338  | 2.56% of 13,186 |
| Foot Dorsal   | 21   | 2.27% of 925    |
| Foot Plantar  | 6    | 5.26% of 114    |
| Toe Dorsal    | 5    | 3.57% of 140    |
| Toe Nail      | 0    | 0% of 8         |
| Toe Plantar   | 1    | 4.17% of 24     |
| Palm Or Sole  | 0    | 0% of 0         |
|               |      |                 |

