


SCARD Pool report for 01-01-2020 to 31-12-2020

| Participants | Doctors | 396 | |
|---------------------|-----------------------------|---------|---------|
| | Patients | 59,605 | |
| | | | |
| Specimens | New lesions | 113,647 | 85.36% |
| | Previously biopsied lesions | 19,491 | 14.64% |
| | Total lesions | 133,138 | 100.00% |

Breakdown of new versus previously biopsied specimens



| Percentage of new lesions tested which were malignant | 62.24% |
|--|--------|
| Percentage of Definitively excised lesions that were malignant | 80.04% |
| Lesions tested to find one melanoma (NNT) | 3.97 |
| Percentage of lesions tested for NMSC which were NMSC | 75.81% |
| Ratio of New BCCs : New Melanomas | 9:1 |

Accuracy

Diagnostic sensitivity

| Melanomas | 75.34% of 3,836 |
|-----------|------------------|
| All NMSC | 95.69% of 64,572 |
| BCCs | 90.96% of 34,621 |
| SCCs | 82.77% of 29,597 |
| | |

Positive predictive value

| Melanomas | 41.45% of 6,972 |
|-----------|------------------|
| All NMSC | 81.67% of 75,661 |
| BCCs | 41.62% of 75,661 |
| SCCs | 65.72% of 37,274 |

Adequacy


Percentage of definitively managed malignant cases where surgical margins were adequate

| BCC | 87.77% of 23,205 |
|---------------------------|------------------|
| IEC/Bowens disease | 77.29% of 8,728 |
| SCC | 86.98% of 8,403 |
| Keratoacanthoma | 90.79% of 1,032 |
| Melanoma - in situ | 75.75% of 2,581 |
| Melanoma - invasive | 73.10% of 606 |
| Melanoma - invasive > 1mm | 42.16% of 102 |
| Other malignant | 71.32% of 129 |



Lesion Breakdown Histological Diagnosis

| BCC (unspecified type) | 118 | 0.10% |
|-----------------------------|--------|--------|
| BCC - Superficial | 10,828 | 9.51% |
| BCC - Nodular/Solid | 19,164 | 16.84% |
| BCC - Aggressive | 4,511 | 3.96% |
| IEC/Bowens disease | 16,451 | 14.45% |
| SCC | 11,593 | 10.19% |
| Keratoacanthoma | 1,553 | 1.36% |
| Pinkus Fibroepithelioma | 16 | 0.01% |
| Merkel cell tumour | 13 | 0.01% |
| Other malignant | 160 | 0.14% |
| NMSC Metastasis | 3 | 0.00% |
| Melanoma - in situ | 2,811 | 2.47% |
| Melanoma - invasive | 765 | 0.67% |
| Melanoma - invasive > 1mm | 232 | 0.20% |
| Melanoma - metastasis | 28 | 0.02% |
| MELTUMP | 162 | 0.14% |
| Naevus - other | 6,998 | 6.15% |
| Naevus - dysplastic/Clark | 3,157 | 2.77% |
| Naevus - blue | 395 | 0.35% |
| Naevus - Spitz/Reed | 119 | 0.10% |
| Naevus - Compound | 15 | 0.01% |
| Solar keratosis | 10,506 | 9.23% |
| Solar lentigo | 1,361 | 1.20% |
| Seborrhoeic keratosis | 5,878 | 5.16% |
| Lentigo Simplex | 1 | 0.00% |
| Lichenoid keratosis (LPLK) | 2,153 | 1.89% |
| Dermatofibroma | 830 | 0.73% |
| Sebaceous gland hyperplasia | 272 | 0.24% |
| Benign cyst | 1,789 | 1.57% |
| Other benign | 7,737 | 6.80% |
| Histology Pending | 164 | 0.14% |
| | | |





Procedures

| Definitive | Surgical | Management | used to | exclude | melanoma |
|--------------|----------|------------|---------|---------|---------------|
| Dellillitive | Surgicar | Management | useu to | exclude | IIIEIAIIUIIIA |

| Definitive Surgical Managem | ent used to exclude melanoma | | |
|-----------------------------|---|--------|--------|
| | Ellipse | 2,989 | 87.14% |
| | Flap | 54 | 1.57% |
| | Graft - SSG | 2 | 0.06% |
| | Graft - FTSG | 12 | 0.35% |
| | No Closure | 11 | 0.32% |
| | Shave/Saucerisation | 65 | 1.90% |
| | Curettage & Cautery | 17 | 0.50% |
| | Liquid N2 freeze/thaw | 6 | 0.17% |
| | PDT | 0 | 0% |
| | Imiquimod | 0 | 0% |
| | 5 FU cream | 1 | 0.03% |
| | GP referral | 179 | 5.22% |
| | Specialist referral | 80 | 2.33% |
| | Other | 7 | 0.20% |
| Biopsy used to exclude mela | noma | | |
| | Punch - sample | 677 | 5.47% |
| | Shave - sample | 1,390 | 11.22% |
| | Incisional | 231 | 1.87% |
| | Punch - removal | 2,214 | 17.88% |
| | Shave - removal | 2,762 | 22.30% |
| | Excisional | 5,005 | 40.41% |
| | Curettage | 44 | 0.36% |
| | Other | 61 | 0.49% |
| Breakdown of definitive man | agement procedures for malignant conditions | | |
| | Ellipse | 38,335 | 63.94% |
| | Flap | 5,584 | 9.31% |
| | Graft - SSG | 369 | 0.62% |
| | Graft - FTSG | 1,364 | 2.27% |
| | No Closure | 249 | 0.42% |
| | Shave/Saucerisation | 1,039 | 1.73% |
| | Curettage & Cautery | 6,949 | 11.59% |
| | Liquid N2 freeze/thaw | 707 | 1.18% |
| | PDT | 85 | 0.14% |
| | Imiquimod | 549 | 0.92% |
| | 5 FU cream | 1,170 | 1.95% |
| | GP referral | 931 | 1.55% |
| | Specialist referral | 1,855 | 3.09% |
| | Other | 187 | 0.31% |
| Breakdown of definitive man | agement procedures for benign conditions | | |
| | Ellipse | 5,240 | 75.93% |
| | Flap | 94 | 1.36% |
| | Graft - SSG | 2 | 0.03% |
| | Graft - FTSG | 9 | 0.13% |
| | No Closure | 43 | 0.62% |
| | Shave/Saucerisation | 406 | 5.88% |
| | Liquid N2 freeze/thaw | 220 | 3.19% |
| | 5 FU cream | 117 | 1.70% |
| | GP referral | 83 | 1.20% |
| | Specialist referral | 36 | 0.52% |
| | Other | 100 | 1.45% |
| | | | |



Procedures (continued)


Percentage of procedures/closures that were complex

Complex Closures 11.04% of 48,476 Complex Closures (inc Curettage & Cautery) 9.98% of 53,651

Locations

Breakdown of melanomas by location and percentage melanomas of total lesions at that location

| Nose | 34 | 0.51% of 6,707 |
|---------------|------|-----------------|
| Lip | 5 | 0.29% of 1,750 |
| Ear | 104 | 2.04% of 5,097 |
| Eyelid | 8 | 0.77% of 1,041 |
| Other face | 328 | 1.77% of 18,532 |
| Scalp | 104 | 2.25% of 4,622 |
| Neck | 143 | 2.43% of 5,873 |
| Shoulder | 265 | 5.02% of 5,278 |
| Chest | 185 | 2.72% of 6,791 |
| Abdomen | 100 | 6.64% of 1,507 |
| Genitalia | 1 | 0.63% of 159 |
| Back | 1279 | 8.06% of 15,877 |
| Buttock | 7 | 2.52% of 278 |
| Arm | 439 | 6.56% of 6,694 |
| Forearm | 214 | 2.58% of 8,307 |
| Hand Dorsal | 8 | 0.17% of 4,745 |
| Hand Palmar | 0 | 0% of 32 |
| Finger Dorsal | 0 | 0% of 992 |
| Finger Nail | 1 | 4.35% of 23 |
| Finger Palmar | 0 | 0% of 16 |
| Thigh | 208 | 6.02% of 3,454 |
| Leg | 370 | 2.63% of 14,057 |
| Foot Dorsal | 22 | 1.97% of 1,118 |
| Foot Plantar | 5 | 5.21% of 96 |
| Toe Dorsal | 3 | 1.96% of 153 |
| Toe Nail | 0 | 0% of 15 |
| Toe Plantar | 1 | 4.76% of 21 |
| Palm Or Sole | 0 | 0% of 0 |
| | | |



Upper Limbs - 17.27%