


SCARD Pool report for 01-01-2021 to 31-12-2021

| Participants | Doctors Patients | 321 65,730 | |
|--------------|-----------------------------|---------------|---------|
| Specimens | New lesions | 121,915 | 84.56% |
| | Previously biopsied lesions | 22,268 | 15.44% |
| | Total lesions | 144,183 | 100.00% |

Breakdown of new versus previously biopsied specimens



| Percentage of new lesions tested which were malignant | 63.92% |
|--|--------|
| Percentage of Definitively excised lesions that were malignant | 81.36% |
| Lesions tested to find one melanoma (NNT) | 3.76 |
| Percentage of lesions tested for NMSC which were NMSC | 78.65% |
| Ratio of New BCCs : New Melanomas | 9:1 |

Accuracy

Diagnostic sensitivity

| Melanomas | 76.53% of 4,317 |
|-----------|------------------|
| All NMSC | 96.06% of 70,742 |
| BCCs | 91.79% of 37,923 |
| SCCs | 82.56% of 32,511 |
| | |

Positive predictive value

| Melanomas | 43.95% of 7,518 |
|-----------|------------------|
| All NMSC | 84.69% of 80,241 |
| BCCs | 43.38% of 80,241 |
| SCCs | 67.87% of 39,550 |

Adequacy


Percentage of definitively managed malignant cases where surgical margins were adequate

| BCC | 88.06% of 25,875 |
|---------------------------|------------------|
| IEC/Bowens disease | 80.81% of 8,871 |
| SCC | 86.52% of 8,994 |
| Keratoacanthoma | 90.59% of 1,073 |
| Melanoma - in situ | 76.30% of 2,916 |
| Melanoma - invasive | 70.24% of 662 |
| Melanoma - invasive > 1mm | 42.06% of 107 |
| Other malignant | 69.05% of 126 |



Lesion Breakdown Histological Diagnosis

| 80 | 0.07% |
|--------|---|
| 11,645 | 9.54% |
| 21,708 | 17.78% |
| 4,490 | 3.68% |
| 18,392 | 15.07% |
| 12,589 | 10.31% |
| 1,530 | 1.25% |
| 19 | 0.02% |
| 17 | 0.01% |
| 172 | 0.14% |
| 5 | 0.00% |
| 3,212 | 2.63% |
| 837 | 0.69% |
| 234 | 0.19% |
| 34 | 0.03% |
| 95 | 0.08% |
| 7,276 | 5.96% |
| 3,580 | 2.93% |
| 441 | 0.36% |
| 171 | 0.14% |
| 58 | 0.05% |
| 10,363 | 8.49% |
| 1,490 | 1.22% |
| 5,707 | 4.67% |
| 2 | 0.00% |
| 2,125 | 1.74% |
| 885 | 0.72% |
| 276 | 0.23% |
| 1,957 | 1.60% |
| 7,853 | 6.43% |
| 168 | 0.14% |
| | 11,645 21,708 4,490 18,392 12,589 1,530 19 17 172 5 3,212 837 234 34 95 7,276 3,580 441 171 58 10,363 1,490 5,707 2 2,125 885 276 1,957 7,853 |





| | -) | Report for | SCAND FUUI |
|---------------------------------------|--|------------|------------|
| Procedures | | | |
| Definitive Surgical N | Nanagement used to exclude melanoma | | |
| , , , , , , , , , , , , , , , , , , , | Ellipse | 3,389 | 86.88% |
| | Flap | 51 | 1.31% |
| | Graft - SSG | 0 | 0% |
| | Graft - FTSG | 8 | 0.21% |
| | No Closure | 9 | 0.23% |
| | Shave/Saucerisation | 35 | 0.90% |
| | Curettage & Cautery | 38 | 0.97% |
| | Liquid N2 freeze/thaw | 2 | 0.05% |
| | PDT | 0 | 0% |
| | Imiquimod | 0 | 0% |
| | 5 FU cream | 0 | 0% |
| | GP referral | 228 | 5.84% |
| | Specialist referral | 126 | 3.23% |
| | Other | 3 | 0.08% |
| Biopsy used to exclu | | | |
| . , | Punch - sample | 698 | 5.36% |
| | Shave - sample | 1,604 | 12.31% |
| | Incisional | 232 | 1.78% |
| | Punch - removal | 2,599 | 19.94% |
| | Shave - removal | 2,520 | 19.34% |
| | Excisional | 5,297 | 40.65% |
| | Curettage | 20 | 0.15% |
| | Other | 58 | 0.45% |
| Breakdown of defini | tive management procedures for malignant cor | nditions | |
| | Ellipse | 41,644 | 63.52% |
| | Flap | 5,408 | 8.25% |
| | Graft - SSG | 289 | 0.44% |
| | Graft - FTSG | 1,530 | 2.33% |
| | No Closure | 218 | 0.33% |
| | Shave/Saucerisation | 1,045 | 1.59% |
| | Curettage & Cautery | 8,537 | 13.02% |
| | Liquid N2 freeze/thaw | 564 | 0.86% |
| | PDT | 105 | 0.16% |
| | Imiquimod | 519 | 0.79% |
| | 5 FU cream | 1,309 | 2.00% |
| | GP referral | 1,338 | 2.04% |
| | Specialist referral | 2,244 | 3.42% |
| | Other | 171 | 0.26% |
| Breakdown of defini | tive management procedures for benign condit | ions | |
| | Ellipse | 5,621 | 78.09% |
| | Flap | 81 | 1.13% |
| | Graft - SSG | 5 | 0.07% |
| | Graft - FTSG | 14 | 0.19% |
| | No Closure | 31 | 0.43% |
| | Shave/Saucerisation | 277 | 3.85% |
| | Liquid N2 freeze/thaw | 106 | 1.47% |
| | 5 FU cream | 106 | 1.47% |
| | CD (| 110 | 7 6 40/ |

GP referral

Other

Specialist referral

1.64%

0.79%

1.39%

118

57

100



Procedures (continued)

Percentage of procedures/closures that were complex

| Complex Closures | 9.03% of 33,609 |
|--|-----------------|
| Complex Closures (inc Curettage & Cautery) | 8.08% of 37,572 |

Locations

Breakdown of melanomas by location and percentage melanomas of total lesions at that location

| Nose | 46 | 0.63% of 7,357 |
|---------------|------|-----------------|
| Lip | 5 | 0.28% of 1,818 |
| Ear | 100 | 1.80% of 5,544 |
| Eyelid | 15 | 1.29% of 1,162 |
| Other face | 418 | 2.03% of 20,605 |
| Scalp | 115 | 2.26% of 5,085 |
| Neck | 197 | 3.13% of 6,304 |
| Shoulder | 316 | 5.62% of 5,627 |
| Chest | 210 | 2.95% of 7,110 |
| Abdomen | 95 | 6.12% of 1,552 |
| Genitalia | 2 | 0.96% of 209 |
| Back | 1360 | 7.87% of 17,283 |
| Buttock | 15 | 5.68% of 264 |
| Arm | 460 | 6.30% of 7,303 |
| Forearm | 272 | 3.06% of 8,886 |
| Hand Dorsal | 8 | 0.17% of 4,697 |
| Hand Palmar | 0 | 0% of 30 |
| Finger Dorsal | 3 | 0.38% of 788 |
| Finger Nail | 0 | 0% of 18 |
| Finger Palmar | 0 | 0% of 11 |
| Thigh | 179 | 4.89% of 3,662 |
| Leg | 454 | 3.07% of 14,780 |
| Foot Dorsal | 35 | 3.07% of 1,139 |
| Foot Plantar | 3 | 2.94% of 102 |
| Toe Dorsal | 4 | 1.97% of 203 |
| Toe Nail | 0 | 0% of 15 |
| Toe Plantar | 1 | 9.09% of 11 |
| Palm Or Sole | 0 | 0% of 0 |
| | | |

