


SCARD Pool report for 01-01-2021 to 31-12-2021

| Participants | Doctors | 323 | |
|--------------|-----------------------------|---------|---------|
| | Patients | 66,014 | |
| Specimens | New lesions | 122,287 | 84.55% |
| | Previously biopsied lesions | 22,354 | 15.45% |
| | Total lesions | 144,641 | 100.00% |

Breakdown of new versus previously biopsied specimens



| Percentage of new lesions tested which were malignant | 63.88% |
|--|--------|
| Percentage of Definitively excised lesions that were malignant | 81.29% |
| Lesions tested to find one melanoma (NNT) | 3.75 |
| Percentage of lesions tested for NMSC which were NMSC | 78.63% |
| Ratio of New BCCs : New Melanomas | 9:1 |

Accuracy

Diagnostic sensitivity

| Melanomas | 76.60% of 4,346 |
|-----------|------------------|
| All NMSC | 96.06% of 70,895 |
| BCCs | 91.78% of 38,002 |
| SCCs | 82.58% of 32,584 |
| | |

Positive predictive value

| Melanomas | 44.08% of 7,553 |
|-----------|------------------|
| All NMSC | 84.68% of 80,425 |
| BCCs | 43.37% of 80,425 |
| SCCs | 67.86% of 39,651 |

Adequacy


Percentage of definitively managed malignant cases where surgical margins were adequate

| BCC | 88.08% of 25,932 |
|---------------------------|------------------|
| IEC/Bowens disease | 80.83% of 8,883 |
| SCC | 86.57% of 9,026 |
| Keratoacanthoma | 90.56% of 1,081 |
| Melanoma - in situ | 76.44% of 2,933 |
| Melanoma - invasive | 70.29% of 663 |
| Melanoma - invasive > 1mm | 42.06% of 107 |
| Other malignant | 69.05% of 126 |



Lesion Breakdown Histological Diagnosis

| BCC (unspecified type) | 80 | 0.07% |
|-----------------------------|--------|--------|
| BCC - Superficial | 11,662 | 9.52% |
| BCC - Nodular/Solid | 21,767 | 17.78% |
| BCC - Aggressive | 4,493 | 3.67% |
| IEC/Bowens disease | 18,409 | 15.03% |
| SCC | 12,637 | 10.32% |
| Keratoacanthoma | 1,538 | 1.26% |
| Pinkus Fibroepithelioma | 19 | 0.02% |
| Merkel cell tumour | 17 | 0.01% |
| Other malignant | 173 | 0.14% |
| NMSC Metastasis | 5 | 0.00% |
| Melanoma - in situ | 3,234 | 2.64% |
| Melanoma - invasive | 842 | 0.69% |
| Melanoma - invasive > 1mm | 236 | 0.19% |
| Melanoma - metastasis | 34 | 0.03% |
| MELTUMP | 95 | 0.08% |
| Naevus - other | 7,335 | 5.99% |
| Naevus - dysplastic/Clark | 3,586 | 2.93% |
| Naevus - blue | 442 | 0.36% |
| Naevus - Spitz/Reed | 172 | 0.14% |
| Naevus - Compound | 58 | 0.05% |
| Solar keratosis | 10,390 | 8.48% |
| Solar lentigo | 1,495 | 1.22% |
| Seborrhoeic keratosis | 5,722 | 4.67% |
| Lentigo Simplex | 2 | 0.00% |
| Lichenoid keratosis (LPLK) | 2,136 | 1.74% |
| Dermatofibroma | 887 | 0.72% |
| Sebaceous gland hyperplasia | 277 | 0.23% |
| Benign cyst | 1,967 | 1.61% |
| Other benign | 7,905 | 6.46% |
| Histology Pending | 168 | 0.14% |





Procedures

| Definitive Surgic | al Management used t | o exclude melanoma |
|-------------------|----------------------|------------------------|
| Deminitive Surgic | ai management asea t | o excidue ilicialionia |

| Definitive Surgical Managem | nent used to exclude melanoma | | | |
|---|--|--------|--------|--|
| | Ellipse | 3,400 | 86.73% | |
| | Flap | 51 | 1.30% | |
| | Graft - SSG | 0 | 0% | |
| | Graft - FTSG | 8 | 0.20% | |
| | No Closure | 9 | 0.23% | |
| | Shave/Saucerisation | 39 | 0.99% | |
| | Curettage & Cautery | 38 | 0.97% | |
| | Liquid N2 freeze/thaw | 2 | 0.05% | |
| | PDT | 0 | 0% | |
| | Imiquimod | 0 | 0% | |
| | 5 FU cream | 0 | 0% | |
| | GP referral | 232 | 5.92% | |
| | Specialist referral | 126 | 3.21% | |
| | Other | 3 | 0.08% | |
| Biopsy used to exclude mela | noma | | | |
| | Punch - sample | 701 | 5.35% | |
| | Shave - sample | 1,604 | 12.25% | |
| | Incisional | 232 | 1.77% | |
| | Punch - removal | 2,605 | 19.89% | |
| | Shave - removal | 2,549 | 19.46% | |
| | Excisional | 5,325 | 40.66% | |
| | Curettage | 20 | 0.15% | |
| | Other | 58 | 0.44% | |
| Breakdown of definitive mar | nagement procedures for malignant conditions | | | |
| | Ellipse | 41,748 | 63.52% | |
| | Flap | 5,418 | 8.24% | |
| | Graft - SSG | 289 | 0.44% | |
| | Graft - FTSG | 1,533 | 2.33% | |
| | No Closure | 218 | 0.33% | |
| | Shave/Saucerisation | 1,046 | 1.59% | |
| | Curettage & Cautery | 8,539 | 12.99% | |
| | Liquid N2 freeze/thaw | 566 | 0.86% | |
| | PDT | 105 | 0.16% | |
| | Imiquimod | 531 | 0.81% | |
| | 5 FU cream | 1,313 | 2.00% | |
| | GP referral | 1,341 | 2.04% | |
| | Specialist referral | 2,259 | 3.44% | |
| | Other | 173 | 0.26% | |
| Breakdown of definitive management procedures for benign conditions | | | | |
| | Ellipse | 5,677 | 77.93% | |
| | Flap | 81 | 1.11% | |
| | Graft - SSG | 5 | 0.07% | |
| | Graft - FTSG | 14 | 0.19% | |
| | No Closure | 31 | 0.43% | |
| | Shave/Saucerisation | 302 | 4.15% | |
| | Liquid N2 freeze/thaw | 107 | 1.47% | |
| | 5 FU cream | 107 | 1.47% | |
| | GP referral | 120 | 1.65% | |
| | Specialist referral | 58 | 0.80% | |
| | Other | 100 | 1.37% | |
| | | | | |

Procedures (continued)


Percentage of procedures/closures that were complex

| Complex Closures | 0.96% of 10,411 |
|--|-----------------|
| Complex Closures (inc Curettage & Cautery) | 0.90% of 11,062 |

Locations

Breakdown of melanomas by location and percentage melanomas of total lesions at that location

| · · · · · · · · · · · · · · · · · · · | | |
|---------------------------------------|------|-----------------|
| Nose | 46 | 0.62% of 7,369 |
| Lip | 5 | 0.27% of 1,824 |
| Ear | 100 | 1.80% of 5,566 |
| Eyelid | 15 | 1.29% of 1,164 |
| Other face | 419 | 2.03% of 20,658 |
| Scalp | 116 | 2.27% of 5,112 |
| Neck | 200 | 3.16% of 6,326 |
| Shoulder | 320 | 5.67% of 5,645 |
| Chest | 211 | 2.96% of 7,133 |
| Abdomen | 95 | 6.09% of 1,559 |
| Genitalia | 2 | 0.96% of 209 |
| Back | 1371 | 7.90% of 17,347 |
| Buttock | 15 | 5.68% of 264 |
| Arm | 462 | 6.31% of 7,325 |
| Forearm | 274 | 3.08% of 8,898 |
| Hand Dorsal | 8 | 0.17% of 4,711 |
| Hand Palmar | 0 | 0% of 30 |
| Finger Dorsal | 3 | 0.38% of 789 |
| Finger Nail | 0 | 0% of 18 |
| Finger Palmar | 0 | 0% of 11 |
| Thigh | 179 | 4.87% of 3,674 |
| Leg | 458 | 3.09% of 14,831 |
| Foot Dorsal | 35 | 3.07% of 1,140 |
| Foot Plantar | 3 | 2.88% of 104 |
| Toe Dorsal | 4 | 1.96% of 204 |
| Toe Nail | 0 | 0% of 15 |
| Toe Plantar | 1 | 9.09% of 11 |
| Palm Or Sole | 0 | 0% of 0 |
| | | |



Upper Limbs - 17.20%